

Poster:

Study of early screening of potato (*Solanum tuberosum* L.) as a result of drought stress (*in vitro*)

¹Charloq, ²Ernitha Panjaitan, and ³Biliter A. Sirait

¹Department of Agriculture, Sumatera Utara University, Medan 20155, Indonesia; ²Department of Agriculture, Methodist Indonesia University, Medan 20132, Indonesia; ³Department of Agriculture, Kopertis Wil.I employed of Darma Agung University Medan 20153, Indonesia. Coresponding Author: charloq@yahoo.com

Abstract. Preliminary studies in Screening of Potato (*Solanum tuberosum* L.) as a result of drought stress (*in vitro*). This study aim was to determine the limits of potato (*Solanum tuberosum* L.) to drought stress conditions *in vitro* using PEG. This research was conducted at the Tissue Culture Laboratory Unit / Laboratorium Kultur Jaringan UPT/Balai Benih Induk Hortikultura Dinas Pertanian Propinsi Sumatera Utara, Jl. Karya Jasa No. 6 Gedung Johor Medan in March 2012 to June 2012. The Completely Randomized Design (non factorial) was used with PEG concentrations of five levels, namely: P1 = 10%, P2 = 20%, P3= 30%, P4 = 40% and P5 = 50% with 5 replications. The results showed that increasing the concentration of PEG will reduce the percentage of plantlets survival, reduce plant height, number of roots and number of nodes. Treatment of 50% PEG resulted in the death of potato cuttings due to drought stress.

Keywords: early studies, screening, drought stress

Introduction

Potatoes are one of the alternative commodities that support government programs for food security. In recent years, the need for potatoes is likely to increase and this plant is one of the priority commodity development. Potatoes in Indonesia is used as a source of carbohydrate in the food diversification program. Polyethylene Glycol (PEG) was reported able to hold water so that it becomes not available to plants. The amount of PEG solution to hold water depends on the molecular weight and concentration (Mexal *et al.*, 1975), it is soluble in water, not toxic to plants, and is not easily absorbed making PEG as an effective compound to simulate drought conditions (Mullahey *et al.*, 1996, Dami & Hughes, 1997, Kaur *et al.* 1998). Selection methods were developed with a solution of PEG to identify potato plants respond to drought stress *in vitro*, thus might also consistent *ex vitro*. Van Sint January *et al.* (1997) also argues that it is the consensus that drought tolerant plants is also mediated at the cellular level.

Materials and Methods

The experiment was conducted at the Tissue Culture Laboratory Unit Laboratorium Kultur Jaringan UPT/Balai Benih Induk Hortikultura Dinas Pertanian Propinsi Sumatera Utara, Jl. Karya Jasa No. 6 Gedung Johor Medan in March 2012 to June 2012. This study used the basic media of Murashige and Skoog (MS). The study uses CRD (completely randomized design) non factorial was used with PEG treatment of 5 level which repeated 5 times, namely: P1 = 10%, P2 = 20%, P3= 30%, P4 = 40% and P5= 50%. Granola potato explants prepared by growing potatoes in a polybags containing soil and manure mixture. After 25 days, young potato used as a source of explants. Explants were obtained by excising the young plant potatoes, then planted on liquid MS medium with the addition of PEG 6000 at a concentration of 10%, 20%, 30%, 40 % and 50% (Figure 1). The parameters observed at the end of the study were percentage of living plantlets (%), plant height (cm), number of roots, and number of nodes.

Results and Discussion

Percentage of Survive Plantlets. Concentration of PEG was highly significantly reducing the percentage of survival plantlets (Figure 2). The relationship between the concentration of PEG to the percentage of survival explants follow negative linear regression with an equation of $Y = -0.666x + 112$, $R^2 = 0.726$, the higher the concentration of the PEG, the lower the percentage of living plantlets, that means that potato cuttings are sensitive to PEG treatment *in vitro* and was not able to survive much longer.

Plant height. The relationship between the concentration of PEG with potato explants plant height at 6 weeks age was also follow the negative linear regression with an equation of $Y = -0.075x + 5.242$, $R^2 = 0.902$ (Figure 3). In this case, the potential decline in water media by

Poster:

the addition of PEG, which in turn lead to the decreased of proliferation of tissue explants, growth and shoot regeneration. The Number of Roots. The relationship between the concentration of PEG with the number of roots of potato explants also following the negative linear regression, with an equation of $Y = -0.062x + 4.112$, $R^2 = 0.939$ (Figure 4). The Number of Nodes. The relationship between the concentration of PEG with the number of nodes was following the linear regression with an equation of $Y = -0.068x + 5.105$, $R^2 = -0.904$ (Figure 5). The use of PEG as osmoticum was suspected to stress can reduce plant cell elongation and expansion, which in turn lowered the growth rate. Sirait, *et al.* (2012) also showed that PEG treatment significantly lowered the number of roots, number of nodes, number of leaves, plant height and wet weight of explant.



Figure 1. Planlet Potatoes Growing *in vitro* at the ages of Six Weeks with PEG Treatment




Figure 2. Relationship Concentration of PEG with Survival Plantlets Percentages




Figure 3. Relationship Concentration of PEG with Plant height

Poster:




Figure 4. Relationship Concentration of PEG with Number of Roots




Figure 5. Relationship Concentration of PEG with Number of Nodes

Conclusions

Conclusion. Increasing the concentration of PEG reduce the percentage of plantlets survival, reduce plant height, number of roots and number of nodes *in vitro*. Suggestions. Drought Tolerant (*in vitro*) needs to be evaluated in physiological character and molecular testing.

References

- Asih K., Karjadi, Buchory A. 2007. Micropropagation and production of mini tubers 15 clones introduced from the International Potato Center (CIP-PERU). Proceedings of the National Seminar on Potato Week, Lembang 20 to 21 August 2008. Department of Agricultural Research and Development Research Center and Horticulture Development.
- Bao JS., Yang CS., Xue JQ., Hao YC., 1991. The effect of water stress during different growth periods of maize on its physiological characteristics. *Acta Agronomica Sinica*, 17 (4) : 261-266.
- Bates LS., Waldren RP., Teare ID. 1973. Rapid determination of free proline for water stress studies. *Plant and Soil* 39: 205-207.
- Mexal J., Fisher JT., Osteryoung J., Patrick Reid CP. 1975. Oxygen availability in polyethylene glycol solutions and its implications in plant-water relation. *Plant Physiol* 55:20-24.
- Mullahey JJ., West SH., Cornell JA. 1996. Effects of simulated drought by polyethylene glycol on bahia grass germination. *Seed Sci Technol* 24:219-224 in Wahyu Widoretno, Edi Guhardja, Satriyas Ilyas, Sudarsono, 2001. Effectiveness of Polyethylene Glycol to evaluate the response of soybean genotypes to drought stress at germination phase.
- Sirait, Samosir, Sitingjak, 2012. Effect of GA3 and PEG on growth potato (*Solanum tuberosum*, L) *in vitro*. Departemen of Agriculture, Darma Agung University Medan.
- Van Sint Jan, V., C. C. de Macedo, J. M. Kinet, Bouharmont J. 1997. Selection of Al-resistant plants from a sensitive rice cultivar, using somaclonal variation, *in vitro* and hydroponic cultures. *Euphytica* 97: 303-310.